



Machine Learning

KMA Solaiman
UMBC CMSC 478

Today:

- Bayes Classifiers
- Conditional Independence
- Naïve Bayes

Slide credit: Tom M. Mitchell. - Thanks!

Two Principles for Estimating Parameters

- Maximum Likelihood Estimate (MLE): choose θ that maximizes probability of observed data \mathcal{D}

$$\hat{\theta} = \arg \max_{\theta} P(\mathcal{D} | \theta)$$

- Maximum a Posteriori (MAP) estimate: choose θ that is most probable given prior probability and the data

$$\begin{aligned} \hat{\theta} &= \arg \max_{\theta} P(\theta | \mathcal{D}) \\ &= \arg \max_{\theta} = \frac{P(\mathcal{D} | \theta)P(\theta)}{P(\mathcal{D})} \end{aligned}$$

Maximum Likelihood Estimate



$X=1$ $X=0$
 $P(X=1) = \theta$
 $P(X=0) = 1-\theta$
(Bernoulli)

- Each flip yields boolean value for X

$$X \sim \text{Bernoulli}: P(X) = \theta^X (1 - \theta)^{(1-X)}$$

- Data set D of independent, identically distributed (iid) flips produces α_1 ones, α_0 zeros

$$P(D|\theta) = P(\alpha_1, \alpha_0|\theta) = \theta^{\alpha_1} (1 - \theta)^{\alpha_0}$$

$$\hat{\theta}^{MLE} = \arg \max_{\theta} P(D|\theta) = \frac{\alpha_1}{\alpha_1 + \alpha_0}$$

Maximum A Posteriori (MAP) Estimate



- Data set D of independent, identically distributed (iid) flips produces α_1 ones, α_0 zeros

$$P(D|\theta) = P(\alpha_1, \alpha_0|\theta) = \theta^{\alpha_1}(1 - \theta)^{\alpha_0}$$


- Assume prior $P(\theta) = \text{Beta}(\beta_1, \beta_0) = \frac{1}{B(\beta_1, \beta_0)} \theta^{\beta_1-1}(1 - \theta)^{\beta_0-1}$
- Then

$$\hat{\theta}^{MAP} = \arg \max_{\theta} P(D|\theta)P(\theta) = \frac{\alpha_1 + \beta_1 - 1}{(\alpha_1 + \beta_1 - 1) + (\alpha_0 + \beta_0 - 1)}$$

(like MLE, but hallucinating $\beta_1 - 1$ additional heads, $\beta_0 - 1$ additional tails)

Let's learn classifiers by learning $P(Y|X)$

Consider $Y=Wealth$, $X=<Gender, HoursWorked>$



Gender	HrsWorked	P(rich G,HW)	P(poor G,HW)
F	<40.5	.09	.91
F	>40.5	.21	.79
M	<40.5	.23	.77
M	>40.5	.38	.62

How many parameters must we estimate?

Suppose $X = \langle X_1, \dots, X_n \rangle$

where X_i and Y are boolean RV's

Gender	HrsWorked	P(rich G,HW)	P(poor G,HW)
F	<40.5	.09	.91
F	>40.5	.21	.79
M	<40.5	.23	.77
M	>40.5	.38	.62

To estimate $P(Y | X_1, X_2, \dots, X_n)$

If we have 30 boolean X_i 's: $P(Y | X_1, X_2, \dots, X_{30})$

Bayes Rule

$$P(Y|X) = \frac{P(X|Y)P(Y)}{P(X)}$$

Which is shorthand for:

$$(\forall i, j) P(Y = y_i | X = x_j) = \frac{P(X = x_j | Y = y_i) P(Y = y_i)}{P(X = x_j)}$$

Equivalently:

$$(\forall i, j) P(Y = y_i | X = x_j) = \frac{P(X = x_j | Y = y_i) P(Y = y_i)}{\sum_k P(X = x_j | Y = y_k) P(Y = y_k)}$$

Can we reduce params using Bayes Rule?

Suppose $X = \langle X_1, \dots, X_n \rangle$

where X_i and Y are boolean RV's

$$P(Y|X) = \frac{P(X|Y)P(Y)}{P(X)}$$

How many parameters to define $P(X_1, \dots, X_n | Y)$?

How many parameters to define $P(Y)$?

Can we reduce params using Bayes Rule?

Suppose $X = \langle X_1, \dots, X_n \rangle$

where X_i and Y are boolean RV's

$$P(Y|X) = \frac{P(X|Y)P(Y)}{P(X)}$$

How many parameters to define $P(X_1, \dots, X_n | Y)$?

$$P(X|Y=1) \text{ ----- } 2^n - 1$$

$$P(X|Y=0) \text{ ----- } 2^n - 1$$

How many parameters to define $P(Y)$?

Can we reduce params using Bayes Rule?

Suppose $X = \langle X_1, \dots, X_n \rangle$

where X_i and Y are boolean RV's

$$P(Y|X) = \frac{P(X|Y)P(Y)}{P(X)}$$

how many params for $P(X_1 \dots X_n | Y)$ $(2^n - 1) \cdot 2$

how many for $P(Y) = 1$

Naïve Bayes

Naïve Bayes assumes

$$P(X_1 \dots X_n | Y) = \prod_i P(X_i | Y)$$

i.e., that X_i and X_j are conditionally independent given Y , for all $i \neq j$

Conditional Independence

Definition: X is conditionally independent of Y given Z , if the probability distribution governing X is independent of the value of Y , given the value of Z

$$(\forall i, j, k) P(X = x_i | Y = y_j, Z = z_k) = P(X = x_i | Z = z_k)$$

Which we often write

$$P(X|Y, Z) = P(X|Z)$$

E.g.,

$$P(\textit{Thunder} | \textit{Rain}, \textit{Lightning}) = P(\textit{Thunder} | \textit{Lightning})$$

Naïve Bayes uses assumption that the X_i are conditionally independent, given Y . E.g., $P(X_1|X_2, Y) = P(X_1|Y)$

Given this assumption, then:

$$P(X_1, X_2|Y) =$$

Naïve Bayes uses assumption that the X_i are conditionally independent, given Y . E.g., $P(X_1|X_2, Y) = P(X_1|Y)$

Given this assumption, then:

$$\begin{aligned} P(X_1, X_2|Y) &= P(X_1|X_2, Y)P(X_2|Y) \\ &= P(X_1|Y)P(X_2|Y) \end{aligned}$$

in general: $P(X_1 \dots X_n|Y) = \prod_i P(X_i|Y)$

Naïve Bayes uses assumption that the X_i are conditionally independent, given Y . E.g., $P(X_1|X_2, Y) = P(X_1|Y)$

Given this assumption, then:

$$\begin{aligned} P(X_1, X_2|Y) &= P(X_1|X_2, Y)P(X_2|Y) && \text{Chain Rule} \\ &= P(X_1|Y)P(X_2|Y) \end{aligned}$$

in general:
$$P(X_1 \dots X_n|Y) = \prod_i P(X_i|Y)$$

Naïve Bayes uses assumption that the X_i are conditionally independent, given Y . E.g., $P(X_1|X_2, Y) = P(X_1|Y)$

Given this assumption, then:

$$\begin{aligned} P(X_1, X_2|Y) &= P(X_1|X_2, Y)P(X_2|Y) && \text{Chain Rule} \\ &= P(X_1|Y)P(X_2|Y) \end{aligned}$$

in general:
$$P(X_1 \dots X_n|Y) = \prod_i P(X_i|Y)$$

How many parameters to describe $P(X_1 \dots X_n|Y)$? $P(Y)$?

- Without conditional indep assumption?
- With conditional indep assumption?

Naïve Bayes uses assumption that the X_i are conditionally independent, given Y

Given this assumption, then:

$$\begin{aligned} P(X_1, X_2|Y) &= P(X_1|X_2, Y)P(X_2|Y) \\ &= P(X_1|Y)P(X_2|Y) \end{aligned}$$

Chain rule
Cond. Indep.

in general:
$$P(X_1 \dots X_n|Y) = \prod_i P(X_i|Y)$$

How many parameters to describe $P(X_1 \dots X_n|Y)$? $P(Y)$?

- Without conditional indep assumption? $2(2^n - 1) + 1$
- With conditional indep assumption? $2n + 1$

Naïve Bayes in a Nutshell

Bayes rule:

$$P(Y = y_k | X_1 \dots X_n) = \frac{P(Y = y_k) P(X_1 \dots X_n | Y = y_k)}{\sum_j P(Y = y_j) P(X_1 \dots X_n | Y = y_j)}$$

Assuming conditional independence among X_i 's:

$$P(Y = y_k | X_1 \dots X_n) = \frac{P(Y = y_k) \prod_i P(X_i | Y = y_k)}{\sum_j P(Y = y_j) \prod_i P(X_i | Y = y_j)}$$

So, to pick most probable Y for $X^{new} = \langle X_1, \dots, X_n \rangle$

$$Y^{new} \leftarrow \arg \max_{y_k} P(Y = y_k) \prod_i P(X_i^{new} | Y = y_k)$$

Naïve Bayes Algorithm – discrete X_i

- Train Naïve Bayes (examples)

for each* value y_k

estimate $\pi_k \equiv P(Y = y_k)$

for each* value x_{ij} of each attribute X_i

estimate $\theta_{ijk} \equiv P(X_i = x_{ij} | Y = y_k)$

- Classify (X^{new})

$$Y^{new} \leftarrow \arg \max_{y_k} P(Y = y_k) \prod_i P(X_i^{new} | Y = y_k)$$

$$Y^{new} \leftarrow \arg \max_{y_k} \pi_k \prod_i \theta_{ijk}$$

* probabilities must sum to 1, so need estimate only n-1 of these...

Estimating Parameters: Y, X_i discrete-valued

Maximum likelihood estimates (MLE' s):

$$\hat{\pi}_k = \hat{P}(Y = y_k) = \frac{\#D\{Y = y_k\}}{|D|}$$

$$\hat{\theta}_{ijk} = \hat{P}(X_i = x_{ij} | Y = y_k) = \frac{\#D\{X_i = x_{ij} \wedge Y = y_k\}}{\#D\{Y = y_k\}}$$

Number of items in
dataset D for which $Y=y_k$

Example: Live in Sq Hill? $P(S|G,D,M)$

80, 8

- $S=1$ iff live in Squirrel Hill
- $G=1$ iff shop at SH Giant Eagle
- $D=1$ iff Drive to CMU
- $M=1$ iff Rachel Maddow fan

What probability parameters must we estimate?

Example: Live in Sq Hill? $P(S|G,D,M)$

80, 8

- $S=1$ iff live in Squirrel Hill
- $G=1$ iff shop at SH Giant Eagle
- $D=1$ iff Drive to CMU
- $M=1$ iff Rachel Maddow fan

What probability parameters must we estimate?

$P(S=1)$:

$P(D=1 | S=1)$:

$P(D=1 | S=0)$:

$P(G=1 | S=1)$:

$P(G=1 | S=0)$:

$P(M=1 | S=1)$:

$P(M=1 | S=0)$:

$P(S=0)$:

$P(D=0 | S=1)$:

$P(D=0 | S=0)$:

$P(G=0 | S=1)$:

$P(G=0 | S=0)$:

$P(M=0 | S=1)$:

$P(M=0 | S=0)$:

Example: Live in Sq Hill? $P(S|G,D,B)$

- $S=1$ iff live in Squirrel Hill
- $G=1$ iff shop at SH Giant Eagle
- $D=1$ iff Drive or carpool to CMU
- $B=1$ iff Birthday is before July 1

What probability parameters must we estimate?

Example: Live in Sq Hill? $P(S|G,D,E)$

- $S=1$ iff live in Squirrel Hill
- $G=1$ iff shop at SH Giant Eagle
- $D=1$ iff Drive or Carpool to CMU
- $B=1$ iff Birthday is before July 1

$P(S=1)$:

$P(D=1 | S=1)$:

$P(D=1 | S=0)$:

$P(G=1 | S=1)$:

$P(G=1 | S=0)$:

$P(B=1 | S=1)$:

$P(B=1 | S=0)$:

$P(S=0)$:

$P(D=0 | S=1)$:

$P(D=0 | S=0)$:

$P(G=0 | S=1)$:

$P(G=0 | S=0)$:

$P(B=0 | S=1)$:

$P(B=0 | S=0)$:

Naïve Bayes: Subtlety #1

Often the X_i are not really conditionally independent

- We use Naïve Bayes in many cases anyway, and it often works pretty well
 - often the right classification, even when not the right probability (see [Domingos&Pazzani, 1996])
- What is effect on estimated $P(Y|X)$?
 - Extreme case: what if we add two copies: $X_i = X_k$

Extreme case: what if we add two copies: $X_i = X_k$

Extreme case: what if we add two copies: $X_i = X_k$

$$P(Y=y|X) \propto P(Y=y) \prod_i P(X_i=x | Y=y)$$

$\underbrace{\hspace{10em}}_{P(X_1, \dots, X_n | Y=y)}$

Naïve Bayes: Subtlety #2

If unlucky, our MLE estimate for $P(X_i | Y)$ might be zero.
(for example, $X_i = \text{birthdate}$. $X_i = \text{Jan_25_1992}$)

- Why worry about just one parameter out of many?
- What can be done to address this?

Naïve Bayes: Subtlety #2

If unlucky, our MLE estimate for $P(X_i | Y)$ might be zero. (e.g., $X_i = \text{Birthday_Is_January_30_1992}$)

- Why worry about just one parameter out of many?

$$P(Y|X) \propto P(Y) \prod_i P(X_i = x^{\text{New}} | Y)$$

0

- What can be done to address this?

Estimating Parameters

- Maximum Likelihood Estimate (MLE): choose θ that maximizes probability of observed data \mathcal{D}

$$\hat{\theta} = \arg \max_{\theta} P(\mathcal{D} | \theta)$$

- Maximum a Posteriori (MAP) estimate: choose θ that is most probable given prior probability and the data

$$\begin{aligned} \hat{\theta} &= \arg \max_{\theta} P(\theta | \mathcal{D}) \\ &= \arg \max_{\theta} \frac{P(\mathcal{D} | \theta)P(\theta)}{P(\mathcal{D})} \end{aligned}$$

Estimating Parameters: Y, X_i discrete-valued

Maximum likelihood estimates:

$$\hat{\pi}_k = \hat{P}(Y = y_k) = \frac{\#D\{Y = y_k\}}{|D|}$$

$$\hat{\theta}_{ijk} = \hat{P}(X_i = x_j | Y = y_k) = \frac{\#D\{X_i = x_j \wedge Y = y_k\}}{\#D\{Y = y_k\}}$$

MAP estimates (Beta, Dirichlet priors):

$$\hat{\pi}_k = \hat{P}(Y = y_k) = \frac{\#D\{Y = y_k\} + (\beta_k - 1)}{|D| + \sum_m (\beta_m - 1)}$$

Only difference:
“imaginary” examples

$$\hat{\theta}_{ijk} = \hat{P}(X_i = x_j | Y = y_k) = \frac{\#D\{X_i = x_j \wedge Y = y_k\} + (\beta_k - 1)}{\#D\{Y = y_k\} + \sum_m (\beta_m - 1)}$$

Learning to classify text documents

- Classify which emails are spam?
- Classify which emails promise an attachment?
- Classify which web pages are student home pages?

How shall we represent text documents for Naïve Bayes?

Baseline: Bag of Words Approach

the world of

TOTAL



all about the company

Our energy exploration, production, and distribution operations span the globe, with activities in more than 100 countries.

At TOTAL, we draw our greatest strength from our fast-growing oil and gas reserves. Our strategic emphasis on natural gas provides a strong position in a rapidly expanding market.

Our expanding refining and marketing operations in Asia and the Mediterranean Rim complement already solid positions in Europe, Africa, and the U.S.

Our growing specialty chemicals sector adds balance and profit to the core energy business.

► All About The Company

- Global Activities
- Corporate Structure
- TOTAL's Story
- Upstream Strategy
- Downstream Strategy
- Chemicals Strategy
- TOTAL Foundation
- Homepage



aardvark	0
about	2
all	2
Africa	1
apple	0
anxious	0
...	
gas	1
...	
oil	1
...	
Zaire	0

Learning to classify document: $P(Y|X)$ the “Bag of Words” model

- Y discrete valued. e.g., Spam or not
- $X = \langle X_1, X_2, \dots, X_n \rangle =$ document
- X_i is a random variable describing the word at position i in the document
- possible values for X_i : any word w_k in English
- Document = bag of words: the vector of counts for all w_k 's
 - like #heads, #tails, but we have many more than 2 values
 - assume word probabilities are position independent (i.i.d. rolls of a 50,000-sided die)

Naïve Bayes Algorithm – discrete X_i

- Train Naïve Bayes (examples)

for each value y_k

estimate $\pi_k \equiv P(Y = y_k)$

for each value x_j of each attribute X_i

estimate $\theta_{ijk} \equiv P(X_i = x_j | Y = y_k)$

prob that word x_j appears
in position i , given $Y=y_k$

- Classify (X^{new})

$$Y^{new} \leftarrow \arg \max_{y_k} P(Y = y_k) \prod_i P(X_i^{new} | Y = y_k)$$

$$Y^{new} \leftarrow \arg \max_{y_k} \pi_k \prod_i \theta_{ijk}$$

* Additional assumption: word probabilities are position independent

$$\theta_{ijk} = \theta_{mjk} \text{ for all } i, m$$